

United Nations Educational Scientific and Cultural Organization
and
International Atomic Energy Agency
THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**COMPACTNESS THEOREMS
FOR COUPLED YANG-MILLS FIELDS**

Xi Zhang¹

*Department of Mathematics, Zhejiang University,
Hangzhou 310027, Zhejiang, People's Republic of China*

and

The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

Abstract

In this paper, we consider Coupled Yang-Mills fields on vector bundle E over compact Riemannian manifold M . Under appropriate conditions on the curvature and the Higgs field, two compactness theorems are proved.

MIRAMARE – TRIESTE

July 2004

¹mzhangxi@ictp.trieste.it; xizhang@zju.edu.cn

1 Introduction

Let M be a n -dimensional manifold with a Riemannian metric g , and E a vector bundle over M with a compact Lie group G as its structure group. A connection A of E can be given by specifying a covariant derivative

$$D_A : C^\infty(E) \rightarrow C^\infty(E \otimes \Omega^1 M).$$

In local trivialization of E , D_A is of the form $d + \alpha$ for some $\text{Lie}(G)$ -valued 1-form α . The curvature of A is a $\text{Lie}(G)$ -valued 2-form F_A , which is equal to D_A^2 . As usual, it measures deviation from the symmetry of second derivatives. Suppose that the bundle E has a Riemannian structure. All connections here are required to be compatible with the Riemannian structure.

The Determinant of the volume bundle over M is a line bundle of conformal weight n . We denote by L , the determinant bundle raised to the $\frac{1}{n}$ power. Sections of this bundle are constant in a fixed coordinate system but have weight 1 under scale transformations. Let the Higgs field Φ be a section of $E \otimes L$, and the mass m be a section of L . With these definitions, the Yang-Mills-Higgs functional is defined in the following

$$YMH(A, \Phi) = \int_M [|F_A|^2 + |\nabla_A \Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2] \quad (1.1)$$

where λ is a constant. The Yang-Mills-Higgs equations are

$$\begin{cases} D_A^* F_A = -D_A \Phi \otimes \Phi^*, \\ D_A^* D_A \Phi = -\frac{\lambda}{2}(|\Phi|^2 - m^2)\Phi \end{cases} \quad (1.2)$$

which are the Euler-Lagrange equations of the Yang-Mills-Higgs functional and generalize the Yang-Mills equations. One pair (A, Φ) satisfying the Yang-Mills-Higgs equations will be called by one Coupled Yang-Mills fields.

Gauge transformation σ is a section of $\text{Aut}(E)$ which acts on connections, curvature forms, and Higgs fields according to the transformations:

$$\begin{aligned} \sigma^*(A) &= \sigma^{-1} d\sigma + \sigma^{-1} A \sigma, \\ F_{\sigma^* A} &= \sigma^{-1} F_A \sigma, \\ \sigma^*(\Phi) &= \sigma \circ \Phi. \end{aligned} \quad (1.3)$$

The pair (A, Φ) is gauge equivalent to $(\bar{A}, \bar{\Phi})$ if there is a gauge transformation σ such that $\bar{A} = \sigma^*(A)$ and $\bar{\Phi} = \sigma^*(\Phi)$. It is obvious that the Yang-Mills-Higgs equations are invariant under gauge transformations.

In analytical aspect of the Yang-Mills theory, two most fundamental results are the K. Uhlenbeck's compactness theorem and removable singularity theorem ([Uh1],[Uh2]). Later, the removable singularity theorem of K.Uhlenbeck was extended to coupled Yang-Mills fields by T.H.Parker, L.M.Sibner and R.J.Sibner, P.D.Smith, T.H.Otway ([Pa1], [SS], [Si], [Sm], [OS], [O]).

The modulo space of Yang-Mills connections (or coupled Yang-Mills fields) is the quotient of the set of solutions of the Yang-Mills equation (Yang-Mills-Higgs equations with fixed λ and m) by the gauge group, which consists of all gauge transformations. It is well-known that this modulo space may not be compact. Given any sequence of Yang-Mills connections $\{A_i\}$ with a uniformly bounded L^2 -norm of curvature, Uhlenbeck ([Uh1]) (also see [Na], [Ti]) proved that by taking a subsequence if necessary, A_i converges to, modulo gauge transformations, a Yang-Mills connection A in the smooth topology outside a closed subset $S_b(\{A_i\})$ of Hausdorff codimension at least 4. If M is a 4-dimensional compact manifold, the blow-up locus consists of finitely many points, and the limiting connection A can be extended to be a Yang-Mills connection on the whole manifold with smaller L^2 -norm of curvature ([Uh1], [Uh2]). With M of higher dimension, G.Tian [Ti] studied the geometric structures of the blow-up loci of Yang-Mills connections. He also proved a removable singularity theorem for stationary Yang-Mills connections. Particularly, this implies that the limiting connection A extends to be become a smooth connection on $M \setminus S$ for a closed subset S with vanishing $(n - 4)$ -dimensional Hausdorff measure $H^{n-4}(S) = 0$.

In this paper we consider the compactness property of sequences of coupled Yang-Mills fields (A_i, Φ_i) with a uniformly bounded Yang-Mills-Higgs energy $YMH(A_i, \phi_i)$. We first derive a monotonicity formula of Coupled Yang-Mills fields. Using R.Schoen and K.Uhlenbeck's argument in [Sc] for harmonic maps (the method has been extended to pure Yang-Mills fields by H.Nakajima [Na]), we can deduce ϵ estimates for Coupled Yang-Mills fields. Then, discussing like that in [Ti] (for pure Yang-Mills fields case), we can obtain the following compactness theorem.

Theorem 1.1 *Let (A_i, Φ_i) be a sequence of smooth Coupled Yang-Mills fields over M with fixed $\lambda > 0$ and m ; then there is a subsequence (A_i, Φ_i) and gauge transformations σ_k , such that $\sigma_k^*(A_k, \Phi_k)$ converges to a Coupled Yang-Mills fields (A, Φ) outside a closed subset S of Hausdorff codimension at least 4.*

In the second part of this paper, we want to deduce another compactness theorem under different assumption. Noting that the Yang-Mills-Higgs energy is not conformally ("scale") invariant unless the dimension of underground Riemannian manifolds are four. In higher dimension, only $L^{\frac{n}{4}}$ norm of the Yang-Mills-Higgs energy density is conformally invariant. Under a uniform bound on $L^{\frac{n}{4}}$ norm of the Yang-Mills-Higgs energy density, we deduce the following theorem.

Theorem 1.2 *Let E be a vector bundle over compact Riemannian manifold M of dimension $n > 4$, (A_i, Φ_i) be a sequence of smooth Coupled Yang-Mills fields on E with fixed $\lambda > 0$ and m . Assume that*

$$\int_M [|F_{A_i}|^2 + |\nabla_{A_i} \Phi_i|^2 + \frac{\lambda}{4} (|\Phi_i|^2 - m^2)^2]^{\frac{n}{4}} \leq C_0, \quad (1.4)$$

where C_0 is a positive constant, then there is a subsequence (A_α, Φ_α) converge to, modulo gauge transformations, a smooth Coupled Yang-Mills field (A, Φ) in C^∞ -topology on M .

The ideal of the proof of theorem 1.2 is similar as that in [Zh] for pure Yang-Mills connection case. First, We will use the local curvature estimate and the conformally invariant of $L^{\frac{n}{4}}$ to deduce that there exists a subsequence (A_k, Φ_k) (modulo gauge transformations) converge to a Coupled Yang-Mills connection (A_∞, Φ_∞) in smooth topology outside a blow up set \tilde{S} which is at most finite points. Secondly, using the removable singularity theorem due to L.M.Sibner[Si], we know that the limiting field can be extends to a smooth one on M . Furthermore, discussing as that in [Ti], we will construct non-trivial bubbling fields on R^m as (A_k, Φ_k) approach to (A_∞, Φ_∞) if the blow up set is non-empty. On the other hand, when $n > 4$, we can prove an non-existence theorem for coupled Yang-Mills fields which will show that bubbling fields, in fact, are not exist, then the blow up set \tilde{S} must be empty. So the subsequence (A_k, Φ_k) (modulo gauge transformations) converge to a smooth Coupled Yang-Mills field in C^∞ topology on M .

2 Preliminary Results

Let $\pi : E \rightarrow M$ is a vector bundle over a Riemannian manifold (M, g) with a compact Lie group G as its structure group. A connection A on E is defined by specifying a covariant derivative

$$D = D_A : C^\infty(E) \rightarrow C^\infty(E \otimes \Omega^1 M).$$

where $C^\infty(E)$ denotes the space of C^∞ sections of the bundle E . In a local trivialization $(U_\alpha, \varphi_\alpha)$ of E , the covariant derivative takes the form

$$D = d + A_\alpha, A_\alpha : U_\alpha \rightarrow T^*U_\alpha \otimes \mathfrak{g}$$

Denote the Lie algebra of G by \mathfrak{g} and the adjoint and automorphism bundles by AdE and $AutE$. Assume also E has a metric compatible with the action of G and an inclusion $G \subset SO(r)$. We use the metric on G induced by the trace inner product metric on $SO(r)$.

The gauge group is $\mathbf{G} = C^\infty(AutE)$ which consists of all smooth sections of the bundle $AutE$. Any σ in \mathbf{G} is called a gauge transformation. Two smooth connections A_1 and A_2 of E are equivalent if there is a gauge transformation σ such that $A_2 = \sigma(A_1)$, where $\sigma(A)$ be the connection with $D_{\sigma(A)} = \sigma \cdot D_A \cdot \sigma^{-1}$. One can easily show $\sigma(A) = \sigma \cdot A \cdot \sigma^{-1} - d\sigma \cdot \sigma^{-1}$.

For any connection A of E , its curvature F_A measures the extent to which covariant derivative fail to commute. Then $F_A = D_A^2$ is a section of $\wedge^2 T^*M \otimes AdE$. In each local trivialization $(U_\alpha, \varphi_\alpha)$, we have

$$F_A = dA_\alpha + A_\alpha \wedge A_\alpha. \tag{2.1}$$

The Determinant of the volume bundle over M is a line bundle of conformal weight n . We denote by L , the determinant bundle raised to the $\frac{1}{n}$ power. Sections of this bundle are constant in a fixed coordinate system but have weight 1 under scale transformations. The Higgs field Φ is a section of $E \otimes L$. The mass m is defined to be a section of L , and hence, constant in a fixed coordinate system, but having weight 1 under scale changes. (For a careful and rigorous discussion of conformal weights, see Parker [Pa1], [Pa2]). From these, we construct the Yang-Mills-Higgs Energy with these definitions, the Yang-Mills-Higgs functional is defined in the following

$$YMH(A, \Phi) = \int_M [|F_A|^2 + |\nabla_A \Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2] dv_g. \quad (2.2)$$

where λ is a constant (in this paper, we only consider the case $\lambda > 0$), and the norms are taken with respect to Riemannian metric g and fixed invariant inner products on the bundles AdE , L and E .

Coupled Yang-Mills fields (A, Φ) are the critical points for the above Yang-Mills-Higgs functional. Equivalently, the pairs (A, Φ) satisfy the following equations.

$$\begin{cases} D_A^* F_A = -D_A \Phi \otimes \Phi^*, \\ D_A^* D_A \Phi = -\frac{\lambda}{2}(|\Phi|^2 - m^2)\Phi \end{cases} \quad (2.3)$$

where Φ^* is the adjoint of Φ taken with respect to the inner product of E , and D_A^* denotes the adjoint operator of D_A with respect the Killing form of \mathfrak{g} and the Riemannian metric g on M . The above equations also be called by Yang-Mills-Higgs equations. It is obvious that the Yang-Mills-Higgs energy is invariant by the gauge transformation. Then, if (A, Φ) is a Coupled Yang-Mills field, so is $\sigma^*(A, \Phi)$ for any gauge transformation σ . In other words, the Yang-Mills-Higgs equations are invariant under the action of the gauge group. From the conformal weights of Higgs fields, one can easily show that the Yang-Mills-Higgs equations are also invariant under re-scaling.

Next, we want to deduce a monotonicity formula of the Yang-Mills-Higgs energy, the discussion is similar as that in [P] (also [Ti]) for pure Yang-Mills connections. Let $\{f_t\}_{|t|<\infty}$ be a one-parameter family of diffeomorphisms of M , and A_0 be a fixed smooth connection of E and D be its associated covariant derivative. Then for any connection A , we can define a family of connections $A^t = f_t^*(A)$ as follows: Denote by τ_t^0 the parallel transport of E associate to A_0 along the path $f_s(x)_{0 \leq s \leq t}$, here $x \in M$. We define $A^t = f_t^*(A)$ by defining its associated covariant derivative

$$D_X^t v = (\tau_t^0)^{-1}(D_{df_t(X)} \tau_t^0(v))$$

for any $X \in TM$, $v \in \Gamma(E)$; and define $\Phi^t = f_t^* \Phi(x) = (\tau_t^0)^{-1}(\Phi(f_t(x)))$.

Let X be the vector field $\frac{\partial f_t}{\partial t}|_{t=0}$, by direct calculation, we have the following equality.

$$\begin{aligned} \frac{d}{dt} YMH(A^t, \Phi^t)|_{t=0} = & - \int_M (|F_A|^2 \text{div} X - 4 \sum \langle F_A(\nabla_{e_i} X, e_j), F_A(e_i, e_j) \rangle) dv_g \\ & - \int_M (|\nabla_A \Phi|^2 \text{div} X - \sum 2 \langle \nabla_{\nabla_{e_i} X} \Phi, \nabla_{e_i} \Phi \rangle) dv_g \\ & - \frac{\lambda}{4} \int_M (|\Phi|^2 - m^2)^2 \text{div} X dv_g \end{aligned} \quad (2.4)$$

Fixed any $p \in M$, let r_p be a positive number with properties: there are normal coordinates x_1, \dots, x_n in the geodesic ball $B_p(R_p)$ of (M, g) , such that $p = (0, \dots, 0)$ and for some positive constant $c(p)$,

$$|g_{ij} - \delta_{ij}| \leq c(p)(|x_1|^2 + \dots + |x_n|^2),$$

$$|dg_{ij}| \leq c(p)\sqrt{|x_1|^2 + \dots + |x_n|^2},$$

where $g_{ij} = g(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_j})$. The constant r_p and $c(p)$ can be chosen depending only on the injective radius at p and the curvature of (M, g) . If M is the Euclidean space, we can take $r_p = \infty$ and $c(p) = 0$.

Let $r(x)$ be the distance function from p , and ϕ be a positive function on the unit sphere S^{n-1} . Define vector field

$$X(x) = \xi(r)\phi(\frac{x}{r})r\frac{\partial}{\partial r},$$

where ξ is smooth function with compact support in $B_p(r_p)$. Let $\{e_1, \dots, e_n\}$ be any orthonormal basis near p such that $e_1 = \frac{\partial}{\partial r}$. We have

$$\nabla_{\frac{\partial}{\partial r}} \frac{\partial}{\partial r} = 0,$$

$$\nabla_{\frac{\partial}{\partial r}} X = (\xi' r + \xi)\phi(\theta)\frac{\partial}{\partial r},$$

$$\nabla_{e_i} X = \xi r e_i(\phi)\frac{\partial}{\partial r} + \xi\phi \sum b_{ij} e_j$$

where $|b_{ij} - \delta_{ij}| = O(1)c(p)r^2$, and $i \geq 2$. We will always denote by $O(1)$ a quantity bounded by a constant depending only on n .

Let (A, Φ) be a Coupled Yang-Mills field, applying the above equalities to the first variational formula (2.4), we have

$$\begin{aligned} \int_M |F_A|^2 (\xi' r + (n-4)\xi)\phi + \int_M |\nabla_A \Phi|^2 (\xi' r + (n-2)\xi)\phi + \frac{\lambda}{4} \int_M (|\Phi|^2 - m^2)^2 (\xi' r + n\xi)\phi \\ = - \sum \int_M \xi\phi(b_{ij} - \delta_{ij}) \langle e_j, e_i \rangle (|F_A|^2 + |\nabla_A \Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2) \\ + 4 \sum \int_M \xi\phi(b_{ij} - \delta_{ij}) \langle F_A(e_j, e_k), F_A(e_i, e_k) \rangle \\ + 2 \sum \int_M \xi\phi(b_{ij} - \delta_{ij}) \langle \nabla_{e_j} \Phi, \nabla_{e_i} \Phi \rangle \\ + 4 \int_M \{ \xi' r \phi |F_A(\frac{\partial}{\partial r}, \cdot)|^2 + \xi r \langle F_A(\frac{\partial}{\partial r}, \cdot), F_A(\nabla \phi, \cdot) \rangle \} \\ + 2 \int_M \{ \xi' r \phi |\nabla_{\frac{\partial}{\partial r}} \Phi|^2 + \xi r \langle \nabla_{\frac{\partial}{\partial r}} \Phi, \nabla_{\nabla \phi} \Phi \rangle \} \end{aligned} \quad (2.5)$$

For any τ small enough, we choose $\xi(r) = \xi_\tau(r) = \eta(\frac{r}{\tau})$, where η is smooth and satisfies: $\eta(r) = 1$ for $r \in [0, 1]$, $\eta(r) = 0$ for $r \in [1 + \epsilon, \infty)$, $\epsilon > 0$ and $\eta'(r) \leq 0$. Then

$$\tau \frac{\partial}{\partial \tau} (\xi_\tau(r)) = -r \xi'_\tau(r). \quad (2.6)$$

Plugging this into (2.5), and choosing a number a large enough, we obtain:

$$\begin{aligned} \frac{\partial}{\partial \tau} (\tau^{4-n} e^{a\tau^2} \int_M \xi_\tau \phi |F_A|^2) + \tau^2 \frac{\partial}{\partial \tau} (\tau^{2-n} e^{a\tau^2} \int_M \xi_\tau \phi |\nabla_A \Phi|^2) \\ + \tau^4 \frac{\partial}{\partial \tau} (\frac{\lambda}{4} \tau^{-n} e^{a\tau^2} \int_M \xi_\tau \phi (|\Phi|^2 - m^2)^2) \\ \geq 4\tau^{4-n} e^{a\tau^2} \frac{\partial}{\partial \tau} (\int_M \xi_\tau \phi |F_A(\frac{\partial}{\partial r}, \cdot)|^2) + 2\tau^{4-n} e^{a\tau^2} \frac{\partial}{\partial \tau} (\int_M \xi_\tau \phi |\nabla_{\frac{\partial}{\partial r}} \Phi|^2) \\ - 4\tau^{3-n} e^{a\tau^2} \int_M \xi_\tau r \langle F_A(\frac{\partial}{\partial r}, \cdot), F_A(\nabla \phi, \cdot) \rangle - 2\tau^{3-n} e^{a\tau^2} \int_M \xi_\tau r \langle \nabla_{\frac{\partial}{\partial r}} \Phi, \nabla_{\nabla \phi} \Phi \rangle. \end{aligned} \quad (2.7)$$

Then, by integrating on τ and letting ϵ tend to zero, we have

$$\begin{aligned}
& \rho^{4-n} e^{a\rho^2} \int_{B_p(\rho)} \phi(|F_A|^2 + |\nabla\Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2) dv_g \\
& \quad - \sigma^{4-n} e^{a\sigma^2} \int_{B_p(\sigma)} \phi(|F_A|^2 + |\nabla\Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2) dv_g \\
& \geq - \int_{\sigma}^{\rho} \tau^{3-n} e^{a\tau^2} \int_{B_p(\tau)} r(4|F_A(\frac{\partial}{\partial r}, \cdot)| |F_A(\nabla\phi, \cdot)| + 2|\nabla_{\frac{\partial}{\partial r}}\Phi| |\nabla_{\nabla\phi}\Phi|) dv_g d\tau \\
& \quad + \int_{\sigma}^{\rho} 2\tau^{3-n} e^{a\tau^2} \int_{B_p(\tau)} \phi|\nabla\Phi|^2 dv_g d\tau + \int_{\sigma}^{\rho} 4\tau^{3-n} e^{a\tau^2} \int_{B_p(\tau)} \frac{\lambda}{4}\phi(|\Phi|^2 - m^2)^2 dv_g d\tau \\
& \quad + \int_{B_p(\rho) \subset B_p(\sigma)} r^{4-n} e^{a\tau^2} \phi(4|F_A(\frac{\partial}{\partial r}, \cdot)|^2 + 2|\nabla_{\frac{\partial}{\partial r}}\Phi|^2) dv_g
\end{aligned} \tag{2.8}$$

Taking $\phi \equiv 1$, we obtain the following monotonicity formula for Coupled Yang-Mills fields.

Lemma 2.1(A monotonicity formula): *There exist constants r_P , a depend only on M , such that for any $0 < \rho < \gamma < r_P$, we have*

$$\begin{aligned}
& \rho^{4-n} e^{a\rho^2} \int_{B_p(\rho)} (|F_A|^2 + |\nabla\Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2) dv_g \\
& \quad - \sigma^{4-n} e^{a\sigma^2} \int_{B_p(\sigma)} (|F_A|^2 + |\nabla\Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2) dv_g \\
& \geq \int_{B_p(\rho) \subset B_p(\sigma)} r^{4-n} e^{a\tau^2} (4|F_A(\frac{\partial}{\partial r}, \cdot)|^2 + 2|\nabla_{\frac{\partial}{\partial r}}\Phi|^2) dv_g \\
& \quad + \int_{\sigma}^{\rho} 2\tau^{3-n} e^{a\tau^2} \int_{B_p(\tau)} |\nabla\Phi|^2 dv_g d\tau + \int_{\sigma}^{\rho} 4\tau^{3-n} e^{a\tau^2} \int_{B_p(\tau)} \frac{\lambda}{4}(|\Phi|^2 - m^2)^2 dv_g d\tau.
\end{aligned} \tag{2.9}$$

Moreover, if $M = R^m$ and g is flat, then the equality holds in (2.9) for $\rho \in (0, \infty)$ and $a = 0$.

In the following, we give a ϵ -estimate for Coupled Yang-Mills fields. This estimate was first derived by K.Uhlenbeck [Uh1] (also see [Na], [Ti]) for the curvature of Yang-Mills connections. For convenience, we will denote

$$h = h(A, \Phi) = \sqrt{|F_A|^2 + |\nabla_A\Phi|^2 + \frac{\lambda}{4}(|\Phi|^2 - m^2)^2}. \tag{2.10}$$

Theorem 2.2: *Let (A, Φ) be a coupled Yang-Mills field on a G -bundle E over M . Then there are ϵ and constant C_0 , which depend only on manifold M , such that for any $p \in M$ and $\rho < r_p$, whenever*

$$\rho^{4-n} \int_{B_p(\rho)} h^2 dv_g \leq \epsilon, \tag{2.11}$$

then

$$h(p) \leq \frac{C}{\rho^2} (\rho^{4-n} \int_{B_p(\rho)} h^2 dv_g)^{\frac{1}{2}}. \tag{2.12}$$

Proof. Using Weitzenbock formula, and Yang-Mills-Higgs equation, we have

$$\begin{aligned}
\Delta|F_A|^2 &= 2|\nabla_A F_A|^2 + 2\langle \nabla_A^2 F_A, F_A \rangle \\
&= 2|\nabla_A F_A|^2 - 2\langle (D_A D_A^* F_A + R_M \sharp F_A + F_A \sharp F_A), F_A \rangle \\
&\geq 2|\nabla_A F_A|^2 - 2|F_A|^2(|R_M| + |F_M|) - 2|F_A|(|D_A\Phi|^2 + |\Phi|^2|F_A|),
\end{aligned} \tag{2.13}$$

$$\begin{aligned}
\Delta|D_A\Phi|^2 &= -2 < (D_A^*D_A + D_AD_A^*)D_A\Phi + R_M\sharp D_A\Phi + F_A\sharp D_A\Phi, D_A\Phi > \\
&\quad + 2|\nabla(D_A\Phi)|^2 \\
&= 2|\nabla D_A\Phi|^2 - 2 < R_M\sharp D_A\Phi + F_A\sharp D_A\Phi, D_A\Phi > - 2 < (D_A^*F_A)\Phi, D_A\Phi > \\
&\quad + < *(F_A \wedge D_A\Phi), D_A\Phi > + \lambda(|\Phi|^2 - m^2)|D_A\Phi|^2 + 2\lambda| < D_A\Phi, \Phi > |^2 \\
&\geq 2|\nabla D_A\Phi|^2 - 2(|R_M|^2 + 4|F_A| + 2|\Phi|^2)|\nabla A\Phi|^2 \\
&\quad - \lambda||\Phi|^2 - m^2||D_A\Phi|^2 + 2\lambda| < D_A\Phi, \Phi > |^2,
\end{aligned} \tag{2.14}$$

$$\begin{aligned}
\Delta(|\Phi|^2 - m^2)^2 &= 2|\nabla(|\Phi|^2 - m^2)|^2 + 4(|\Phi|^2 - m^2)|D_A\Phi|^2 \\
&\quad - 4(|\Phi|^2 - m^2) < D_A^*D_A\Phi + F_A\sharp\Phi, \Phi > \\
&\geq 2|\nabla(|\Phi|^2 - m^2)|^2 + 2\lambda(|\Phi|^2 - m^2)^2|\Phi|^2 \\
&\quad - 4||\Phi|^2 - m^2||D_A\Phi|^2 - 4||\Phi|^2 - m^2||F_A||\Phi|^2,
\end{aligned} \tag{2.15}$$

where R_M denote the Riemannian curvature tensor of M . In (), we have used the following equality,

$$D_A^*D_AD_A\Phi = D_A^*(F_A\Phi) = (D_A^*F_A)\Phi - *(F_A \wedge D_A\Phi). \tag{2.16}$$

From above inequalities, we have

$$\begin{aligned}
\Delta h^2 &= 2|\nabla h|^2 + 2h\Delta h \\
&\geq 2|\nabla F_A|^2 + 2|\nabla_A(D_A\Phi)|^2 + \frac{\lambda}{2}|\nabla(|\Phi|^2 - m^2)|^2 \\
&\quad - (4m^2 + 8|R_M|)h^2 - (14 + 4\lambda^{-\frac{1}{2}} + 2\lambda^{\frac{1}{2}})h^3.
\end{aligned} \tag{2.17}$$

On the other hand, it is easy to check that

$$|\nabla h|^2 \leq |\nabla|F_A||^2 + |\nabla|D_A\Phi||^2 + \frac{\lambda}{4}|D(|\Phi|^2 - m^2)|^2. \tag{2.18}$$

So we have

$$\Delta h \geq -(4m^2 + 8|R_M|)h - (14 + 4\lambda^{-\frac{1}{2}} + 2\lambda^{\frac{1}{2}})h^2. \tag{2.19}$$

Define a function

$$f(r) = (\rho - 2r)^2 \sup_{x \in B_p(r)} h(x), \tag{2.20}$$

$r \in [0, \frac{1}{2}\rho]$. Then f is continuous in $[0, \frac{1}{2}\rho]$ with $f(\frac{1}{2}\rho) = 0$, so that f attains its maximum at a certain r_0 in $[0, \frac{1}{2}\rho]$.

First we claim that $f(r_0) \leq 64$ if ϵ is sufficiently small. Assume that $f(r_0) > 64$. Put $b = \sup_{x \in B_p(r_0)} h = h(x_0)$, and taking $\sigma = \frac{1}{4}(\rho - 2r_0)$, then we get

$$\begin{aligned}
\sup_{B_{x_0}(\sigma)} h &\leq \sup_{x \in B_p(r_0 + \sigma)} h(x) \\
&\leq \frac{(\rho - 2r_0)^2}{(\rho - 2r_0 - 2\sigma)^2} \sup_{x \in B_p(r_0)} h(x) = 4b.
\end{aligned} \tag{2.21}$$

Clearly, $\sigma\sqrt{b} \geq 2$. Define a metric $g' = bg$, and denote the Yang-Mills-Higgs energy density with respect to g' by $h_{g'}$, then $h_{g'} = b^{-1}h_g$. Hence,

$$\sup_{x \in B_2(x_0, g')} h_{g'} \leq 4, \tag{2.22}$$

where $B_2(x_0, g')$ denotes the geodesic ball of g' with radius 2 and center at x_0 .

Since (A, Φ) is a Coupled Yang-Mills field, and the Yang-Mills-Higgs equations are invariant under scaling, from (2.19), we have

$$\Delta_{g'} h_{g'} \geq -C_1 h_{g'} - C_2 h_{g'}^2, \tag{2.23}$$

where C_1, C_2 are constants depending only on r_p, m, λ , and the sectional curvature bound of (M, g) . Using (2.22), we have

$$\Delta_{g'} h_{g'} \geq -(C_1 + 4C_2) h_{g'}. \quad (2.24)$$

Then, by using either the mean-value theorem or the standard Moser iteration, we have

$$1 = h_{g'}(x_0) \leq C_3 \left(\int_{B_1(x_0, g')} h_{g'}^2 dv_{g'} \right)^{\frac{1}{2}}, \quad (2.25)$$

where C_3 is some uniform constant.

On the other hand, by the monotonicity formula (2.9),

$$\begin{aligned} \int_{B_1(x_0, g')} h_{g'}^2 dv_{g'} &= (\sqrt{b})^{n-4} \int_{B_{x_0}(\frac{1}{\sqrt{b}})} h^2 dv_g \\ &\leq \left(\frac{1}{2}\rho\right)^{4-n} e^{\frac{a}{4}\rho^2} \int_{B_{x_0}(\frac{1}{2}\rho)} h^2 dv_g \\ &\leq 2^{n-4} e^{\frac{a}{4}\rho^2} \epsilon. \end{aligned} \quad (2.26)$$

Combining this with (2.25), we obtain

$$1 \leq C_3 (2^{n-4} e^{\frac{a}{4}\rho^2} \epsilon)^{\frac{1}{2}}. \quad (2.27)$$

But, it is impossible when ϵ is sufficiently small. Then, the claim is proved.

Thus, we have

$$\sup_{x \in B_p(\frac{1}{4}\rho)} \rho^2 h \leq 4f(r_0) \leq 256.$$

Scaling the metric again, let $\tilde{g} = \rho^{-2}g$, then $h_{\tilde{g}} = \rho^2 h_g$. It follows from the above inequality and (2.24) with g' replaced by g that for some uniform constant C_4 ,

$$\Delta_{\tilde{g}} h_{\tilde{g}} \geq -C_4 h_{\tilde{g}}. \quad (2.28)$$

Then (2.12) follows from (2.28) and a standard Moser iteration.

□

Using Hölder inequality, we have the following corollary.

Corollary 2.3: *Let (A, Φ) be a Coupled Yang-Mills field on a G -bundle E over M . Then there are ϵ and constant C_5 , which depend only on manifold M , such that for any $p \in M$ and $\rho < r_p$, whenever*

$$\int_{B_p(\rho)} h^{\frac{n}{2}} dv_g \leq \epsilon, \quad (2.29)$$

then

$$h(p) \leq \frac{C_5}{\rho^2} \left(\int_{B_p(\rho)} h^{\frac{n}{2}} dv_g \right)^{\frac{2}{n}}. \quad (2.30)$$

Using the monotonicity formula (2.9), one can easily prove (theorem 3.3 [Zh]) the following corollary.

Corollary 2.4: *If (A, Φ) is a Coupled Yang-Mills field over (R^n, g_0) ($n \geq 5$) and satisfying*

$$\int_{R^n} h(A, \Phi)^{\frac{n}{2}} dx < \infty,$$

where g_0 is a flat metric, then $h(A, \Phi) \equiv 0$.

In order to compactify the modulo space of Coupled Yang-Mills fields, we need to use singular Coupled Yang-Mills fields of a certain type. An admissible Coupled Yang-Mills field is a smooth field (A, Φ) defined outside a closed subset S in M , such that (1) $H^{n-4}(S \cap K) < \infty$ for any compact subset $K \subset M$, where $H^{n-4}(\cdot)$ stands for the $(n-4)$ -dimensional Hausdorff measure; (2) (A, Φ) satisfies the Yang-Mills-Higgs equations (2.3) $M \setminus S$; (3) (A, Φ) satisfies $\int_{M \setminus S} h^2 dV_g < \infty$. Clearly, (A, Φ) is smooth on M if $S = \emptyset$. We will call S the singular set of coupled field (A, Φ) .

Furthermore, an admissible Coupled Yang-Mills field (A, Φ) is called stationary if (A, Φ) satisfies

$$\begin{aligned} 0 = & - \int_M (|F_A|^2 \operatorname{div} X - 4 \sum \langle F_A(\nabla_{e_i} X, e_j), F_A(e_i, e_j) \rangle) dv_g \\ & - \int_M (|\nabla_A \Phi|^2 \operatorname{div} X - \sum 2 \langle \nabla_{\nabla_{e_i} X} \Phi, \nabla_{e_i} \Phi \rangle) dv_g \\ & - \frac{\lambda}{4} \int_M (|\Phi|^2 - m^2)^2 \operatorname{div} X dv_g \end{aligned} \quad (2.31)$$

for any compactly supported vector field X . Where $\{e_i\}$ is any orthonormal basis of M . If (A, Φ) is a smooth Coupled Yang-Mills field, this follows from the first variation formula (2.4).

Proposition 2.5: *Let $n = \dim M > 4$ and S be a discrete set in M . If (A, Φ) is a Coupled Yang-Mills field on $M \setminus S$ and satisfies $\int_K h^{\frac{m}{2}} dV_g < \infty$ for each compact set $K \subset M$; then (A, Φ) is stationary and the monotonicity formula (2.9) still hold on M .*

Proof: Denote

$$\begin{aligned} \Psi(X) = & - \int_M (|F_A|^2 \operatorname{div} X - 4 \sum \langle F_A(\nabla_{e_i} X, e_j), F_A(e_i, e_j) \rangle) dv_g \\ & - \int_M (|\nabla_A \Phi|^2 \operatorname{div} X - \sum 2 \langle \nabla_{\nabla_{e_i} X} \Phi, \nabla_{e_i} \Phi \rangle) dv_g \\ & - \frac{\lambda}{4} \int_M (|\Phi|^2 - m^2)^2 \operatorname{div} X dv_g, \end{aligned} \quad (2.32)$$

where X is a variation vector field with compact support set and $\{e_i\}$ is an orthonormal frame of TM .

We may assume that S consists of a single point p . For $r > 0$ we take a cut-off function $\eta_r \in C_0^\infty(M)$ satisfying: $0 \leq \eta_r \leq 1$, $|\nabla \eta_r| \leq \frac{2}{r}$ in M and $\eta_r(x) = 1$, if $x \in B_p(r)$; $\eta_r(x) = 0$, if $x \in M \setminus B_p(2r)$. Since (A, Φ) is a Coupled Yang-Mills field on $M \setminus B_p(r)$ for any $r > 0$, we have $\Psi(X - \eta_r X) = 0$ for any $r > 0$. Thus, we have

$$\begin{aligned}
|\Psi(X)| &= |\Psi(\eta_r X)| \leq C \int_M h^2(\eta_r |\nabla X| + |\nabla \eta_r| |X|) dV_g \\
&\leq C(\int_{B_p(2r)} h^2 |\nabla X| dV_g + \frac{1}{r} \int_{B_h(2r)} h^2 |X| dV_g) \\
&\leq C(r^{n-4} \sup_M |\nabla X| + r^{n-5} \sup_M |X|)(\int_{B(P,2r)} h^{\frac{n}{2}} dV_g)^{\frac{4}{n}}.
\end{aligned}$$

By conditions the right-hand side tends to 0 as $r \rightarrow 0$. Hence, we get $\Psi(X) = 0$ for any X . This shows that (A, Φ) is stationary on M . □

3 Proof of theorem 1.1

Suppose that (A_i, Φ_i) be a sequence Coupled Yang-Mills fields on bundle E over Riemannian manifold (M, g) . Let ϵ be sufficiently small and a be as in theorem 2.1. We define a closed subset for each i and $r > 0$ sufficiently small:

$$S_{i,r} = \{x \in M \mid r^{4-n} e^{ar^2} \int_{B_x(r)} h^2 dv_g \geq \epsilon\}. \quad (3.1)$$

It follows from the monotonicity formula (2.9) that $S_{i,r} \subset S_{i,r'}$ for any $r \leq r'$. By the standard diagonal process, we can choose a subsequence which also be denoted by $\{i\}$ such that for each k , the $S_{i,2^{-k}}$ converge to a closed subset $S_{2^{-k}}$. It is obvious that $S_{2^{-k}} \subset S_{2^{-l}}$ for $k \geq l$. Then, put

$$S = \cap_k S_{2^{-k}}. \quad (3.2)$$

Discussing like in [Ti] (Proposition 3.1.2), one can prove that S is of Hausdorff codimension at least 4.

Assume that $x \in M \setminus S$, then by the definition of S , there exist r less than $\text{dist}(x, S)$ and $\text{in} \text{grad}(x)$, and $N = N(x) > 0$, such that for $i \geq N$, we have:

$$r^{4-n} \int_{B_x(r)} h^2(A_i, \Phi_i) dv_g < \epsilon. \quad (3.2)$$

Since the left-hand side of (3.2) is scaling invariant, we may rescale to assume that $r = 4$.

In what follows we shall only consider those $i \geq N$ when we restrict our attention on the geodesic ball $B_x(r)$. The point wise a priori estimate for Coupled Yang-Mills fields (theorem 2.2) and (3.2) imply that there exists a uniform constant $C_5 > 0$ such that

$$\sup_{B_x(1)} h^2(A_i, \Phi_i) \leq C_5 \epsilon, \quad (3.3)$$

if we assume that ϵ is less than that in theorem 2.2. Thus we have

$$\begin{aligned}
\sup_{B_x(1)} |F_{A_i}|^2 + |\nabla_{A_i} \Phi_i|^2 &\leq C_5 \epsilon, \\
\sup_{B_x(1)} |\Phi_i|^2 &\leq C_6
\end{aligned} \quad (3.4)$$

where C_6 is a uniform constant.

Fix a local trivialization for E on $B_x(1)$. If ϵ is sufficiently small, by the first inequality in (3.4), we may apply the result of Uhlenbeck ([Uh1] theorem 2.7) to find Coulomb gauges for A_i on $B_x(1)$. In other words, there exist gauge transformations σ_i such that the connections $A'_i = \sigma_i(A_i)$ satisfy

$$d^* A'_i = 0, \quad \text{on } B_x(1), \quad (3.5)$$

$$d_{\psi}^* A'_{i,\psi} = 0, \quad \text{on } \partial B_x(1), \quad (3.6)$$

$$\|A'_i\|_{L^\infty(B_x(1))} \leq C_7 \|F_{A_i}\|_{L^\infty(B_x(1))} \leq C_8 \sqrt{\epsilon}, \quad (3.7)$$

where $A_{i,\psi}$ denote the restriction of A_i on $\partial B_x(1)$, and C_7, C_8 are uniform constants.

Let $\Phi'_i = \sigma_i \cdot \Phi_i$. The equations (2.3) and (3.4) are gauge invariant and hence they hold if we replace (A_i, Φ_i) by (A'_i, Φ'_i) . We observe that (2.3) and (3.5) form an elliptic system for (A'_i, Φ'_i) over $B_x(1)$. By the standard elliptic theory, we conclude that the bounds on derivatives of (A'_i, Φ'_i) are uniform in i , hence by passing to a subsequence, $\sigma_i^*(A_i, \Phi_i)$ converges to a Coupled Yang-Mills field (A', Φ') in smooth topology on $B_x(\frac{1}{2})$.

We may cover the set $M \setminus S$ by a countable union of balls $B_{x_\alpha}(r_\alpha)$ such that (3.2) apply with $r = 8r_\alpha$ and $x = x_\alpha$. Applying the above analysis to each ball $B_{x_\alpha}(8r_\alpha)$, by passing to a subsequence, there exist smooth gauge transformations $\sigma_{i,\alpha}$, such that the sequence $\sigma_{i,\alpha}^*(A_i, \Phi_i)$ converges in smooth topology to a Coupled Yang-Mills field $(A'_\alpha, \Phi'_\alpha)$ on $B_{x_\alpha}(r_\alpha)$. We can now use a standard diagonal process of gluing gauges ([DK], Theorem 4.4.8), again passing to a subsequence, to obtain smooth gauge transformations σ_i on $M \setminus S$, such that $\sigma_i^*(A_i, \Phi_i)$ converge to a field (A, Φ) in smooth topology on compact subsets of $M \setminus S$. So, we have proved theorem 1.1.

4 Proof of theorem 1.2

Proposition 4.1 ([Si]): *Let (A, Φ) be a stationary Coupled Yang-Mills field on $M \setminus S$, where S is a discrete set. If $\int_K h(A, \Phi)^{\frac{n}{2}} dV_g < \infty$ for each compact set $K \subset M$, where $n > 4$ is the dimension of M , then there exist a gauge transformation σ such that $\sigma^*(A, \Phi)$ can be extended to be a smooth Coupled Yang-Mills field on M .*

Theorem 4.2: *Let $\{(A_i, \Phi_i)\}$ be a sequence of smooth Coupled Yang-Mills fields on E with $\int_M h^{\frac{n}{2}} dv_g \leq \Lambda$; then there exist a subsequence $\{\alpha\} \subset \{i\}$ and a (possibly empty) finite set $\tilde{S} = \{p_k\}_{k=1}^l$ of M satisfying the following:*

(1), *the subsequence (A_α, Φ_α) converge to a smooth Yang-Mills connection (A, Φ) in the C^∞ -topology on $M \setminus \Sigma$.*

(2), for each $k = 1, \dots, l$, there exists constants $\theta_k > 0$ such that

$$h(A_\alpha, \Phi_\alpha)^{\frac{n}{2}} dV_g \longrightarrow h(A, \Phi)^{\frac{n}{2}} dV_g + \sum_{k=1}^l \theta_k \cdot \delta_{p_k} \quad (4.1)$$

weakly in the sense of Radon measures on M . Here δ_{p_k} denotes dirac measure.

Proof: Let ϵ be as in the proof of theorem 1.1. We define a closed subset for each i and $r > 0$;

$$E_{i,r} = \{x \in M \mid \int_{B_r(x)} h(A_i, \Phi_i)^{\frac{n}{2}} dV_g \geq \epsilon\}. \quad (4.2)$$

It is obvious that $E_{i,r} \subset E_{i,R}$ for any $r \leq R$. By the standard diagonal process, we can choose a subsequence $\{i_j\}$ of $\{i\}$ such that for each k , the $E_{i_j, 2^{-k}}$ converge to a closed subset $E_{2^{-k}}$. Then $E_{2^{-k}} \subset E_{2^{-l}}$ for $k \geq l$.

Put $S = \cap_k E_{2^{-k}}$. We first claim that S is at most a finite set. We fixed an arbitrary compact set $K \subset M$. For any $\delta > 0$ sufficiently small, let $\{B_{x_b}(4\delta)\}$ be any finite covering of $S \cap K$ such that $x_b \in S \cap K$; $B_{x_b}(2\delta) \cap B_{x_c}(2\delta) = \emptyset$ for $b \neq c$. Take k big enough such that $2^{-k} < \delta$. Then for j sufficiently large, there are $y_b \in E_{i_j, 2^{-k}}$ such that $d(x_b, y_b) < \delta$. Then $\{B_{y_b}(5\delta)\}$ is a finite covering of $S \cap K$ and $B_{y_b}(\delta) \cap B_{y_c}(\delta) = \emptyset$ for $b \neq c$. On the other hand, for each b

$$\int_{B_{y_b}(\delta)} h(A_{i_j}, \Phi_{i_j})^{\frac{n}{2}} dV_g \geq \epsilon. \quad (4.3)$$

Summing up, we get

$$I \leq \frac{1}{\epsilon} \sum_{b=1}^I \int_{B_{y_b}(\delta)} h(A_{i_j}, \Phi_{i_j})^{\frac{n}{2}} dV_g \leq \frac{\Lambda}{\epsilon}, \quad (4.4)$$

This shows $\mathcal{H}^0(S \cap K) \leq \frac{\Lambda}{\epsilon}$ where \mathcal{H}^0 denotes the 0-dimensional Hausdorff measure on M . Since the 0-dimensional Hausdorff measure coincides with the counting measure, $S \cap K$ is at most finite. So S is at most a finite points set.

On the other hand, using corollary 2.3 and discussing like that in the proof of theorem 1.1, one can prove that there exists a subsequence $\{\tilde{i}\} \subset \{i\}$ and gauge transformations $\sigma(\tilde{i})$, such that $\sigma(\tilde{i})^*(A_{\tilde{i}}, \Phi_{\tilde{i}})$ converge to a Coupled Yang-Mills field (A, Φ) in C^∞ -topology on any compact subset outside S . Using Fatou's lemma, we have

$$\int_M h(A, \Phi)^{\frac{n}{2}} dV_g \leq \liminf_{\tilde{i} \rightarrow \infty} \int_M h(A_{\tilde{i}}, \Phi_{\tilde{i}})^{\frac{n}{2}} dV_g \leq \Lambda. \quad (4.5)$$

By Proposition 4.1, there exists a gauge transformation σ such that $\sigma^*(A, \Phi)$ extends to be a smooth Coupled Yang-Mills field on M .

Define

$$\tilde{S} = \bigcap_{r>0} \{x \in M \mid \liminf_{\tilde{i} \rightarrow \infty} \int_{B(r)} h(A_{\tilde{i}}, \Phi_{\tilde{i}})^{\frac{n}{2}} dV_g \geq \epsilon\}. \quad (4.6)$$

Now we want to show that \tilde{S} is contained in the above S . In fact, for any $x_0 \in M \setminus S$, if r is sufficiently small,

$$\int_{B_{x_0}(r)} h(A, \Phi)^{\frac{n}{2}} dV_g < \frac{\epsilon}{4}.$$

This implies that for \tilde{i} sufficiently large,

$$\int_{B_{x_0}(r)} h(A_{\tilde{i}}, \Phi_{\tilde{i}})^{\frac{n}{2}} dV_g < \frac{\epsilon}{2}.$$

Hence, $x_0 \in M \setminus \tilde{S}$. This shows that $\tilde{S} \subset S$.

Suppose $x_0 \in S \setminus \tilde{S}$; then there is an $r_0 > 0$ such that

$$\int_{B_{x_0}(r_0)} h(i')^{\frac{n}{2}} dV_g < \epsilon$$

for some subsequence $i' \rightarrow \infty$. By Corollary 2.3 ,

$$\sup_{x \in B(x_0, \frac{1}{2}r_0)} h(i') \leq C_9 \cdot r_0^2 \cdot \epsilon^{\frac{2}{m}}$$

for some uniform constant C_9 . This implies that there exists some subsequence of $\{A_{i'}, \Phi_{i'}\}$ (modulo gauge transformations) converge in $B_{x_0}(\frac{1}{2}r_0)$ in the C^∞ topology. Then, there exists a subsequence $\{(A_\alpha, \Phi_\alpha)\}$ and a finite set \tilde{S} such that $\{(A_\alpha, \Phi_\alpha)\}$ (modulo gauge transformations) converges to a Coupled Yang-Mills field (A, Φ) in the C^∞ topology on $M \setminus \tilde{S}$.

Consider the Radon measure $\mu_\alpha = h(\alpha)^{\frac{n}{2}} dV_g$. By taking a subsequence if necessary, we may assume that $\mu_\alpha \rightarrow \mu$ weakly on M as Radon measures. Let us write (by Fatou's lemma)

$$\mu = h(A, \Phi)^{\frac{n}{2}} dV_g + \nu \tag{4.7}$$

for some nonnegative Radon measure ν on M . Since $\{(A_\alpha, \Phi_\alpha)\}$ converges to (A, Φ) in the C^∞ topology on $M \setminus \tilde{S}$, the support of measure ν is contained in the discrete set \tilde{S} . Thus, we have $\nu = \sum_{k=1}^l \theta_k \delta_{p_k}$ for some $\theta_k \geq 0$ where we set $\tilde{S} = \{p_k\}_{k=1}^l$.

We show each θ_k is positive. Fix any p_k . For arbitrarily small $r > 0$, we take a cut-off function $\eta_r \in C_0^\infty(M)$ satisfying $0 \leq \eta_r \leq 1$ in M and $\eta_r(x) = 1$ if $x \in B_{p_k}(r)$; $\eta_r(x) = 0$ if $x \in M \setminus B_{p_k}(2r)$. By definition of \tilde{S} we have

$$\epsilon \leq \liminf_{\alpha \rightarrow \infty} \int_{B_{p_k}(r)} h(\alpha)^{\frac{n}{2}} dV_g \leq \lim_{\alpha \rightarrow \infty} \int_M \eta_r h(\alpha)^{\frac{n}{2}} dV_g \leq \theta_k + \int_{B_{p_k}(2r)} h(A, \Phi)^{\frac{n}{2}} dV_g. \tag{4.8}$$

Letting $r \rightarrow 0$, we obtain $\theta_k \geq \epsilon > 0$. This completes the proof. \square

In the following theorem, we will discuss the bubbling phenomenon of Coupled Yang-Mills fields.

Theorem 4.3: Let $\{(A_\alpha, \Phi_\alpha)\}, \tilde{S}$ be as in Theorem 4.2; and $p \in \tilde{S}$. Then there are linear transformations $\tau_\alpha : T_p M \rightarrow T_p M$ such that a subsequence of $\tau_\alpha^* \exp_p^*(A_\alpha, \Phi_\alpha)$ converges to a smooth Coupled Yang-Mills field (B, Ψ) on a trivial bundle over $(T_p M, g_p)$; and satisfying $h(B, \Psi) \neq 0$ and $\int_{T_p M} h(B, \Psi)^{\frac{n}{2}} dv_{g_p} \leq \theta_p$; where θ_p is determined in Theorem 4.2.

Proof: We take a normal coordinate system $\{x^1, \dots, x^n\}$ on the geodesic ball $B_p(2R)$ centered at p , and assume that $\tilde{S} \cap B_p(2R) = \{p\}$. Let $B(x, r)$ be the open ball in the normal coordinates with center x and radius r and let $B(r) = B(0, r)$. Defining the concentration function

$$Y_\alpha(t) = \sup_{y \in B(R)} \int_{\exp_p(B(y, t))} h(A_\alpha, \Phi_\alpha)^{\frac{n}{2}} dV_g, \quad (4.9)$$

for any $0 \leq t < R$. Each function Y_α is continuous and non-decreasing in t , and $Y_\alpha(0) = 0$. On the other hand, by the definition of \tilde{S} , we have

$$Y_\alpha(R) \geq \int_{B_p(R)} h(A_\alpha, \Phi_\alpha)^{\frac{n}{2}} dV_g \geq \frac{3\epsilon}{4} \quad (4.10)$$

holds for sufficiently large α . By continuity of Y_α , there exist $0 < r_\alpha < R$ and $x_\alpha \in \overline{B(R)}$ such that

$$Y_\alpha(r_\alpha) = \int_{\exp_p(B(x_\alpha, r_\alpha))} h(A_\alpha, \Phi_\alpha)^{\frac{n}{2}} dV_g = \frac{\epsilon}{2}. \quad (4.11)$$

Since the p is the a unique point in $\tilde{S} \cap B_p(2R)$, we obtain $r_\alpha \rightarrow 0$, $x_\alpha \rightarrow 0$, as $\alpha \rightarrow \infty$. Defining linear transformations $\tau_\alpha(x) = x_\alpha + r_\alpha \cdot x$ on $T_p M$. Let $U(\alpha) = B(-\frac{x_\alpha}{r_\alpha}, \frac{2R}{r_\alpha}) \subset T_p M$. It is easy to see that $B(2R) = \sigma_\alpha(U(\alpha))$. Since x_α lies in $B(\frac{R}{2})$ for sufficiently large α , we have $B(\frac{R}{r_\alpha}) \subset U(\alpha)$, which leads to $U(\alpha) \rightarrow T_p M$ as $\alpha \rightarrow \infty$.

We set $B_\alpha = \tau_\alpha^* \exp_p^*(A_\alpha)$, $\Psi_\alpha = \tau_\alpha^* \exp_p^*(\Phi_\alpha)$. We can easily see (B_α, Ψ_α) are Coupled Yang-Mills fields on pull backed bundles over $(U(\alpha), g_\alpha)$, where the metric $g_\alpha = r_\alpha^{-2} \tau_\alpha^* \exp_p^* g$. Note that the based manifolds $(T_p M, g_\alpha)$ converge to $(T_p M, g_p) \cong R^n$ as $\alpha \rightarrow \infty$. By the definition of (B_α, Ψ_α) , x_α , r_α , we have

$$\int_{U(\alpha)} h(B_\alpha, \Psi_\alpha)^{\frac{n}{2}} dV_{g_\alpha} = \int_{B_p(2R)} h(A_\alpha, \Phi_\alpha)^{\frac{n}{2}} dV_g \leq \Lambda, \quad (4.12)$$

and

$$Y_\alpha(r_\alpha) = \int_{B(1)} h(B_\alpha, \Psi_\alpha)^{\frac{n}{2}} dV_{g_\alpha} = \sup_{z \in \tau_\alpha^{-1}(B(R))} \int_{B(z, 1)} h(B_\alpha, \Psi_\alpha)^{\frac{n}{2}} dV_{g_\alpha} = \frac{\epsilon}{2}. \quad (4.13)$$

Since $g_\alpha \rightarrow g_p$ in C^∞ topology as $\alpha \rightarrow \infty$, and the constants ϵ and C_5 in Corollary 2.3 depend only on λ and the bound of sectional curvature of metrics. If we assume that the ϵ in (4.13) is small enough, then we can apply corollary 2.3 for any α . So, we have

$$\sup_{B(z, \frac{1}{2})} h(B_\alpha, \Psi_\alpha) \leq C_{10} \epsilon^{\frac{2}{n}}$$

for any $z \in K$, here C_{10} is a uniform constant and K is an arbitrary compact set in $T_p M$. This implies that there exists a subsequence (B_β, Ψ_β) (modulo gauge transformations) converge to a smooth Coupled Yang-Mills field (B, Ψ) over $(T_p(M), g_p)$. Passing to the limit in (4.13), we have

$$\int_{B(1)} h(B, \Psi)^{\frac{n}{2}} dx = \frac{\epsilon}{2}. \quad (4.14)$$

This shows that $h(B, \Psi) \neq 0$. By Fatou's lemma, we have

$$\begin{aligned} \int_{T_p M} h(B, \Psi)^{\frac{n}{2}} dx &\leq \liminf_{\beta \rightarrow \infty} \int_{U(\beta)} h(B_\beta, \Psi_\beta)^{\frac{n}{2}} dV_{g_\beta}, \\ &\leq \theta_p + \int_{B_p(2R)} h(A, \Phi)^{\frac{n}{2}} dV_g \end{aligned}$$

Letting $R \rightarrow 0$, we have

$$\int_{T_p M} h(B, \Psi)^{\frac{n}{2}} dx \leq \theta_p. \quad (4.15)$$

This completes the proof. □

If the blow up set \tilde{S} , is not empty, from theorem 4.3, we know that there exists a bubbling Coupled Yang-Mills field (B, Ψ) over (R^n, g_0) satisfying $h(B, \Psi) \neq 0$ and

$$\int_{R^n} h(B, \Psi)^{\frac{n}{2}} dx \leq \infty.$$

But, this is contradictory with Corollary 2.4. So, the blow up set \tilde{S} which we constructed in Theorem 4.2 must be empty. Then, the subsequence (A_α, Φ_α) (modulo gauge transformations) converges to a smooth Coupled Yang-Mills field (A, Φ) in the C^∞ -topology on M . This completes the proof of Theorem 1.2.

Acknowledgments. This paper was written while the author was visiting the Abdus Salam International Centre for Theoretic Physics, Trieste, Italy. He wishes to acknowledge the generous support of the Centre.

References

- [DK] S.K.Donaldson, P.B.Kronheimer, The geometry of four manifolds, *The Clarendon Press, Oxford University Press*, 1990.
- [Na] H.Nakajima: Compactness of the moduli space of Yang-Mills connections in higher dimensions. *J.Math.Soc.Japan.***40**, 383-392 (1988).
- [O] T.H.Otway: Higher-order singularities in coupled Yang-Mills-Higgs fields. *Nonlinear Analysis*, **12** , No.3, 239-244 (1990).
- [OS] T.H.Otway and L.M.Sibner: Point singularities of coupled gauge fields with low energy. *Com-muns Math. Phys.* **111**, 275-279 (1987).
- [P] P.Price: A monotonicity formula for Yang-Mills fields. *Manuscripta.Math.***43,no.2**, 131-166 (1983).
- [Pa1] T.H.Parker: Gauge theories on four dimensional Riemannian manifolds. *Communs Math. Phys.* **85**, 563-602 (1982).
- [Pa2] T.H.Parker: Conformal fields and stability. *Math.Z.*,**185**, 305-319 (1984).
- [Sc] R.Schoen, Analytic aspects of the harmonic maps problem. *Math. Sci. Res. Inst. Publ.* **2**, Springer-verlag, New York, 321-358, (1984).
- [Si] L.M.Sibner: The isolated point singularity problem for the coupled Yang-Mills equation in higher dimensions.*Math.Ann.***271**, 125-131 (1985).
- [Sm] P.D.Smith: Removable singularities for Yang-Mills-Higgs equations in two dimensional. *Ann. Inst. H. Poincare Anal. Non Lineaire* , **7**, No. 6, 561-588 (1990).
- [SS] L.M.Sibner and R.J.Sibner: Removable singularities of coupled Yang-Mills fields in R^3 . *Com-muns Math. Phys.***93**, 1-17 (1984).
- [Ti] Tian.G .: Gauge theory and calibrated geometry. *Ann. Math.* **42**, 271-297 (2000).
- [Uh1] K.K.Uhlenbeck, : Removable singularities in Yang-Mills fields. *Comm. Math. Phys* **83**, 11-29 (1982).
- [Uh2] K.K.Uhlenbeck, : Connections with L^p bounds on curvature. *Comm. Math. Phys* **83**, 31-42 (1982).
- [Zh] X.Zhang, A compactness theorem for Yang-Mills connections, *Geometry and nonlinear partial differential equations (Hangzhou 2001)*, *AMS/IP Stud.Adv.Math.*, **29**, 217-225.